

Exact ground states of extended $t - J_z$ models on a square lattice

Zohar Nussinov

Department of Physics, Washington University, St. Louis, MO 63160, USA

Anders Rosengren

Condensed Matter Theory, KTH Physics, AlbaNova University Center, SE-106 91 Stockholm, Sweden

(Dated: Received February 2, 2008)

We examine special extended spin $S = 1/2$ fermionic and hard core bosonic $t - J_z$ models with nearest neighbor and next nearest neighbor interactions to find exact ground states. Some of these models display an exponentially large degeneracy with diagonal stripe like patterns.

PACS numbers: 71.27.+a, 71.28.+d, 77.80.-e

I. INTRODUCTION

It has been nearly two decades since the discovery of the high- T_c superconductors [1]. Although much has been learned since, there is still no satisfactory explanation of what causes the superconductivity. There is widespread agreement that it should be possible to describe the electronic properties of the square CuO_2 lattices by the two-dimensional repulsive Hubbard model. This well known model is given by

$$H = -t \sum_{\langle ij \rangle \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + c_{j\sigma}^\dagger c_{i\sigma}) + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (1)$$

where $c_{i\sigma}^\dagger$ creates an electron on site i with spin σ and j is a nearest neighbor of i . This model contains both the movement of the electrons (hopping) (t , kinetic energy) and the interactions of the electrons if they are on the same site (U , potential energy). The Hubbard model is one of the simplest possible models of interacting electrons. In the large U limit, the model reduces to the well known $t - J$ model

$$H = -t \sum_{\langle ij \rangle, \sigma} (\bar{c}_{i,\sigma}^\dagger \bar{c}_{j,\sigma} + h.c.) + J \sum_{\langle ij \rangle} \vec{s}_i \cdot \vec{s}_j \quad (2)$$

and the now constrained fermion operators $\bar{c}_{i,\sigma}^\dagger, \bar{c}_{i,\sigma}$ obey the constraint of no double occupancy at any site i ($n_i = \sum_\sigma \bar{c}_{i,\sigma}^\dagger \bar{c}_{i,\sigma}$ has expectation values 0 or 1). In formal terms, the Fock space is projected to this sector adhering to these constraints. An even simpler version is afforded by the so-called $t - J_z$ model given by

$$H = -t \sum_{\langle ij \rangle, \sigma} (\bar{c}_{i,\sigma}^\dagger \bar{c}_{j,\sigma} + h.c.) + J_z \sum_{\langle ij \rangle} s_i^z s_j^z \quad (3)$$

wherein the $SU(2)$ spin symmetry of the original Hubbard and $t - J$ models has been lifted.

Numerical calculations on such models with open and cylindrical boundary conditions have found non-uniform “stripe” patterns [2] which have earlier been predicted by approximate solutions to the Hubbard model [3], and derived, very convincingly, by the addition of the strong Coulomb effects present in the cuprates [4]. At the moment, there is no clear consensus as to whether these patterns display the genuine ground state of the bare Hubbard (or $t - J$) Hamiltonian

or amount to a finite-size artifact. Attacking these models has proven not to be an easy task. The Hilbert space of the model is too large for exact numerical solutions. Monte Carlo simulations suffer from the sign problem. The interaction energy U is approximately equal to $8t$, so we are neither in an extremely strong, nor in a weakly interacting limit. As has been emphasized by [4], Coulomb repulsions in the cuprates are poorly screened and universally promote the existence of stripes. Notwithstanding, it would be of interest to attain rigorous results on whether various inhomogeneous electronic patterns can, in principle, be realized in the presence of short-range interactions and hopping alone in some models.

The less ambitious goal of this article is not directly related to the very challenging endeavor of understanding such complicated systems (albeit motivated by it). Rather, we construct exactly solvable high dimensional models related to the original $t - J_z$ models. Inspired by exactly solvable AKLT [5, 6] and in particular Klein models [7], [8], [9], [10], [11], we construct an exactly solvable model whose individual pair building blocks are an extension (containing superconducting pair terms and additional phases for the fermionic version) of the $t - J_z$ approximation of the Hubbard model. The price for attaining this solvable short range model (all interactions involve sites which are, at most, two lattice constants apart) and others is that their results cannot be directly related to the models of more physical interest (such as the plain $t - J_z$ model).

II. CONSTRUCTION OF QUASI-EXACT SOLVABLE $S = 1/2$ FERMIONIC AND HARD CORE BOSONIC SYSTEMS ON A SQUARE LATTICE

In the up and coming, we follow a generalization of the Klein models [7], [8], [9], [10], [11] yet now for fermionic $S = 1/2$ particles on the square lattice. To do so, we first find, much in the spirit of [7], [8], [9], [10], [11] a quasi-exactly solvable $S = 1$ model on a square lattice [12]. We then apply a Jordan Wigner transformation to convert the $S = 1$ problem into a fermionic $S = 1/2$ system. The resulting $S = 1/2$ system is none other than a $t - J_z$ model augmented by superconducting pairing terms and phase factors.

A. Construction of ground states for the $S = 1$ system

First, we generalize the Klein model ([7], [8], [9], [10], [11]) constructions to the problem of $S = 1$ spins on the square lattice whose sites are labeled by i (the locations are $\vec{i} = (i_x, i_y)$ are vectors with i_x and i_y integers). We assume that the lattice has periodic boundary conditions. For clarity, we dispense with the vector sign henceforth. We consider a spin $S = 1$ system whose Hamiltonian can be written as a sum over all plaquettes of the lattice

$$H = \sum_{\square} h_{\square}, \quad (4)$$

with $h_{\square}$ a polynomial of the total spin of the four sites forming a given plaquette

$$\vec{S}_{\square} = \sum_{i \in \square} \vec{S}_i, \quad (5)$$

i.e.

$$h_{\square} = \mathcal{P}(\vec{S}_{\square}^2) = \sum_{n=0}^m a_n \vec{S}_{\square}^{2n}. \quad (6)$$

Numerous polynomials (and hence short-range Hamiltonians) can be engineered to find exactly solvable models. Before doing so let us review the possible four and two particle states of $S = 1$ spins. With four spins in tow, the total spin of any plaquette, $0 \leq S_{\square} \leq 4$. The Hilbert space spanned by four $S = 1$ spins can be written as

$$1 \otimes 1 \otimes 1 \otimes 1 = 4 \oplus (3)^3 \oplus (2)^6 \oplus (1)^6 \oplus (0)^3. \quad (7)$$

Here, the left hand side denote the direct product of the four $S = 1$ spins while the right hand side denotes the decomposition in the total spin basis.

The sum of any two nearest neighbor spins on the lattice ($S_{pair}(S_{pair} + 1) = (\vec{S}_i + \vec{S}_j)^2$) can only attain the three values $S_{pair} = 0, 1, 2$. If any two spins forming a given plaquette are in a singlet state ($S_{pair} = 0$), then the total spin of the plaquette cannot exceed two, $S_{\square} \leq 2$. It follows that if the polynomial $\mathcal{P}$ of Eq.(6) is non-negative definite and equal to zero for $S_{\square} = 0, 1, 2$, then any state $|\psi\rangle$ which is a superposition of dimer coverings of the lattice each of which has (at least) one dimer per each plaquette then

$$|\psi\rangle = \sum_P \alpha_P \prod_{ij \in P} |S_{ij}\rangle, \quad (8)$$

(with such dimer coverings labeled by P) is a ground state of H . The above singlet state is given by

$$|S_{ij}\rangle = \frac{1}{\sqrt{3}}[|m_i = 1, m_j = -1\rangle + |m_i = -1, m_j = 1\rangle - |m_i = 0, m_j = 0\rangle]. \quad (9)$$

Putting all of the pieces together, any state of the form Eqs.(8,9) is a ground state of the Hamiltonian of Eqs. (4, 6)

for any non-negative definite polynomial $\mathcal{P}$ which vanishes for $S_{\square} = 0, 1, 2$. For concreteness, we now explicitly give an example of such a polynomial,

$$h_{\square} = \vec{S}_{\square}^2(\vec{S}_{\square}^2 - 2)(\vec{S}_{\square}^2 - 6). \quad (10)$$

Many other non-negative definite polynomials over all possible plaquette spin values can be written down such that they vanish only for $S_{\square} = 0, 1, 2$. Any of the dimer coverings of the form given by Eqs.(8,9) is a ground state of the Hamiltonian given by Eqs.(4, 10).

B. Geometry of the ground states of the $S = 1$ system

Enumerating all possible ground states of the form of Eq.(8) amounts to the task of finding all possible dimer coverings of the on a square lattice with periodic boundary conditions such that every plaquette hosts, at least, one dimer. The problem of hard core dimer coverings was investigated by [13] wherein the number of fully packed coverings on an infinite lattice was found to scale as $N \simeq \exp[GA/\pi]$ with A the number of sites in the lattice and G Catalan's constant ($G = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} \approx 0.915965$). When the condition of (at least) a single dimer per plaquette is imposed on a square lattice with periodic boundary conditions, we find that there are 2^L (with L the lattice length) collinear diagonal coverings[9]. Along each diagonal, we may decide at our whim, to orient the dimers either vertically or horizontally. As the number of diagonals is L and as there are two such choices for dimer orientations (horizontal/vertical) for each diagonal, the number of individual dimer ground states scales as 2^L . Apart from these 2^L "stripe" states, there are no other dimer coverings such that one dimer appears in each plaquette.

Implementing these geometric constraints on the general ground states of the $S = 1$ system of Eqs.(8,9), the allowed ground states read

$$|\psi\rangle = \sum_{P_{\pm}} \alpha_{P_{\pm}} \prod_{i \in A} |S_{i, i+\hat{\phi}(i_x \pm i_y)}\rangle. \quad (11)$$

The difference between the general dimer states of Eq.(8) and Eq.(11) lies in the explicit form of the allowed partitions P which is discussed below. In Eq.(11), the sites $i \in A$ belong to one sub-lattice (say the one given by $(i_x \pm i_y)$ being even with $i_{x,y}$ the x and y components of i) and the unit vectors $\hat{\phi}(i_x \pm i_y) = \hat{e}_x$ or $-\hat{e}_y$ are random Ising variables. The configurations P_+ are those in which the orientation of the dimers (horizontal or vertical) is fixed along right-tilting diagonals ($i_x - i_y = \text{const}$) whereas in P_- are those in which the orientation of the dimers is fixed along left-tilting diagonals (fixed $i_x + i_y$). As there are 2^L possible sequences in P_+ for the field $\hat{\phi}(i_x \pm i_y)$ along all of the right-tilting diagonals (whose number is L) with a similar number of possible coverings P_- , the set of coverings $\{P_+, P_-\}$ spans $\mathcal{O}(2^{L+1})$ configurations. Similarly, additional dimer coverings where the left/bottom most position of the dimer may span both sub-lattices appear. Various orders can be stabilized by augmenting the Hamiltonian by an additional field favoring this or the

other ground states [9]. A cartoon of such an external field of relative strength g stabilizing the parallel (all $\hat{\phi} = \hat{e}_x$ in Eq.(11)) and zig-zag phase ($\hat{\phi} = \hat{e}_x, \hat{e}_y, \hat{e}_x, \hat{e}_y, \dots$ uniformly on consecutive parallel diagonals) at different values of the external field g is shown in Figure 1. Such a term may be, e.g.,

$$H_g = \left[\sum_{i_x+i_y \equiv 0 \pmod{4}} (\vec{S}_i + \vec{S}_{i+\hat{e}_x})^2 - g \sum_{i_x+i_y \equiv 2 \pmod{4}} \{(\vec{S}_i + \vec{S}_{i+\hat{e}_y})^2 - (\vec{S}_i + \vec{S}_{i+\hat{e}_x})^2\} \right]. \quad (12)$$

Here, the first term of Eq.(12), spans sites i belonging to half of the sites of sub-lattice A (the sites in which the sum of the x and y components of the sites $\vec{i}$ is an integer multiple of four, $i_x + i_y = 4m$ with m an integer). The second term in Eq.(12), spans sites which belong to the other half of the sub-lattice A .




FIG. 1: Reproduced from [9]- two possible dimer coverings of the square lattice (a, b) such that each plaquette hosts one dimer. In the context of the $S = 1$ Hamiltonian discussed here, the ovals denote a singlet state $|S_{ij}\rangle$ of Eq.(9) formed between two nearest neighbor sites i and j . Such coverings can be stabilized by augmenting the Hamiltonian of Eq.(4,6) with additional terms similar to those of Eq.(12). As a function of g a transition occurs between the single diagonal (SD) configuration of (a) and the zigzag (ZD) phase of (b). The shorthand O.P. denotes an order parameter in each of the phases.

This model hosts a form of topological order in some of its ground states which is seen by string correlators. Any ground state

$$|\psi_P\rangle = \prod_{ij \in P} |S_{ij}\rangle, \quad (13)$$

is a direct product of two leg ladders (which are oriented along the diagonals) which are fully occupied by singlet states along

each rung. Correlators involving spins along a line of consecutive singlets in P alone display the same behavior as that of a two leg ladder. As the wave-function decouples into a product of two spin singlet wave-functions along each rung of the ladder, it is readily verified that in the states of Eq.(13), the string correlator

$$\langle S_{1,i}^z S_{2,i}^z \exp[i\pi \sum_{k=i+1}^{j-1} (S_{1,k}^z + S_{2,k}^z)] S_{1,j}^z S_{2,j}^z \rangle = \frac{4}{9}, \quad (14)$$

irrespective of the separation between rungs i and j . Here, the indices 1 and 2 denote the location of the spin on one of the endpoints of the rung. A correlator of the form of Eq.(14) was first introduced by Todo et al. [14].

C. An extended $t - J_z S = 1/2$ fermionic model

We now map our solvable $S = 1$ Hamiltonians and their ground states onto an electronic ($S = 1/2$ fermion) problem. To this end, we employ a Jordan-Wigner transformation mapping the three states of a $S = 1$ spin onto the three states of a constrained $S = 1/2$ fermion (one in which a forbidden double occupancy at any given site is strictly enforced).

Rather explicitly, following [15], the mapping between these constrained $S = 1/2$ fermions and the $S = 1$ spin operators is

$$\begin{aligned} S_j^+ &= \sqrt{2}(\bar{c}_{j\uparrow}^\dagger K_j + K_j^\dagger \bar{c}_{j\downarrow}), \\ S_j^- &= \sqrt{2}(K_j^\dagger \bar{c}_{j\uparrow} + \bar{c}_{j\downarrow}^\dagger K_j), \\ S_j^z &= \bar{c}_{j\uparrow}^\dagger \bar{c}_{j\uparrow} - \bar{c}_{j\downarrow}^\dagger \bar{c}_{j\downarrow} \end{aligned} \quad (15)$$

Here the two dimensional kink operators leading to the correct (fermionic) statistics upon transmutation via Aharonov-Bohm phases by flux attachment [16] are

$$K_j \equiv \exp[i \sum_k \theta_{k,j} \bar{n}_k] \quad (16)$$

with $\bar{n}_k = \bar{c}_{k\uparrow}^\dagger \bar{c}_{k\uparrow} + \bar{c}_{k\downarrow}^\dagger \bar{c}_{k\downarrow}$ the total occupancy of site k and $\theta_{k,j}$ the angle between the line connecting sites k and j and, say, the horizontal x -axis. Here the sum spans all lattice sites k .

Explicitly inserting this into the scalar products forming $h_\square$ in Eqs.(4, 10), we find that in $h_\square$ any former appearance of $(\vec{S}_a \cdot \vec{S}_b)$ in the $S = 1$ problem is now replaced by a $t - J_z$ like pair term augmented by superconducting pieces with phases in tow,

$$\begin{aligned} T_{ab} &\equiv 4s_a^z s_b^z + (\bar{c}_{a\uparrow}^\dagger \bar{c}_{b\uparrow} e^{i\phi_{ab}} + \bar{c}_{a\uparrow}^\dagger \bar{c}_{b\downarrow}^\dagger e^{i\bar{\phi}_{ab}} \\ &\quad + \bar{c}_{a\downarrow} \bar{c}_{b\uparrow} e^{-i\bar{\phi}_{ab}} + \bar{c}_{a\downarrow} \bar{c}_{b\downarrow}^\dagger e^{-i\phi_{ab}} + h.c.). \end{aligned} \quad (17)$$

Here, $\vec{s}_{a,b}$ denote the $S = 1/2$ spin operators of the constrained fermions on sites a and b which belong to the same plaquette, and the phases

$$\begin{aligned} \phi_{ab} &\equiv \sum_c [(\theta_{ac} - \theta_{bc}) \bar{n}_c], \\ \bar{\phi}_{ab} &\equiv \sum_c [(\theta_{ac} + \theta_{bc}) \bar{n}_c] \end{aligned} \quad (18)$$

with the sum performed over all lattice sites c . The operator T_{ab} is an extended variant of the nearest neighbor interaction appearing in Eq.(3) with $J_z = 4t$ having additional terms (both additive and multiplicative).

With this substitution any appearance of $\vec{S}_\square^2$ in the $S = 1$ problem is simply replaced by

$$\vec{S}_\square^2 \rightarrow 2\left(\sum_{(ab)} T_{ab} + 4\right) \quad (19)$$

where (ab) span all six pairs of sites which may be chosen from the four sites comprising the plaquette. The Hamiltonian governing the $S = 1/2$ fermionic system is

$$H = \sum_{\square} \left[\left(\sum_{(ab) \in \square} T_{ab} + 1 \right) \times \left(\sum_{(cd) \in \square} T_{cd} + 3 \right) \left(\sum_{(fg) \in \square} T_{fg} + 4 \right) \right]. \quad (20)$$

This is a particular variant of an extended $t - J_z$ model with fixed couplings.

Similarly, the electronic state ground state of Eq.(20) (derived from $|S_{ij}\rangle$) is

$$|\chi_{ij}\rangle = \frac{1}{\sqrt{3}}(|\uparrow_i\downarrow_j\rangle + |\downarrow_i\uparrow_j\rangle - |0_i0_j\rangle). \quad (21)$$

Here, $|\uparrow_i\rangle(|\downarrow_i\rangle)$ denotes a single electron at site i with spin up (down) while $|0_i\rangle$ corresponds to an empty site at site i . Thus a ground state of a Hamiltonian given by Eq.(20) with the definition of Eq.(17) is

$$|\psi\rangle = \sum_P \alpha_P \prod_{ij \in P} |\chi_{ij}\rangle. \quad (22)$$

In $|\psi\rangle$, at all lattice sites k , the occupancy $\langle n_k \rangle = 2/3$ and the phases $\phi_{ab}, \bar{\phi}_{ab}$ vanish if we replace, in Eq.(18), $\bar{n}_k$ by $\langle \bar{n}_k \rangle$. Eq.(22) is superposition of states with different particle numbers; this is a consequence of the superconducting pair creation/annihilation terms in Eq.(17). The number of dimer coverings (the dimer partitioned denoted by P an example of which is shown in Fig.(1)) is exponential in the perimeter of the system

All stated above is exact. In the following paragraph (and here alone) we depart from the rigorous results derived above and briefly speculate about possible extensions. As within the ground state of Eq.(22) there is no kinetic motion of the electrons- we might speculate that the transmutation statistics might well become irrelevant (as no transmutations occur). If this is the case, $|\psi\rangle$ of Eq.(22), will be an eigen-state of Eq.(20) with all phases $\phi_{ab}, \bar{\phi}_{ab}$ set to zero.

D. Hard Core Bose $S = 1/2$ extended $t - J_z$ models

If we replace the hard core $S = 1/2$ fermions by hard core particles obeying other statistics, we find a generalization of the above. Rather explicitly, if hard core particles given by

$\bar{b}_{\alpha,\sigma}, \bar{b}_{\alpha,\sigma}^\dagger$, with $\sigma = \uparrow, \downarrow$, obey the fractional statistics of a phase change of $e^{i\pi\alpha}$ under the interchange of two particles (e.g. $\alpha = 0$ is the hard core bosonic case) then Eq.(22) is a ground state of Eq.(20) subject to the substitution $T_{ab} \rightarrow T_{ab}^\alpha$ with

$$T_{ab}^\alpha \equiv 4s_a^z s_b^z + (\bar{c}_{a\downarrow}^\dagger \bar{c}_{b\uparrow} e^{i\alpha\phi_{ab}} + \bar{c}_{a\uparrow}^\dagger \bar{c}_{b\downarrow} e^{i\alpha\bar{\phi}_{ab}} + \bar{c}_{a\downarrow} \bar{c}_{b\uparrow} e^{-i\alpha\bar{\phi}_{ab}} + \bar{c}_{a\uparrow} \bar{c}_{b\downarrow} e^{-i\alpha\phi_{ab}} + h.c.). \quad (23)$$

In the case of spin $S = 1/2$ hard core bosons [by which we refer to two flavor ($\sigma = \uparrow, \downarrow$) hard core bosons], the resulting Hamiltonian explicitly reads

$$H = \sum_{\square} \left[\left(\sum_{(ab) \in \square} 4s_a^z s_b^z + (\bar{c}_{a\downarrow}^\dagger \bar{c}_{b\uparrow} + \bar{c}_{a\uparrow}^\dagger \bar{c}_{b\downarrow} + \bar{c}_{a\downarrow} \bar{c}_{b\uparrow} + \bar{c}_{a\uparrow} \bar{c}_{b\downarrow} + h.c.) + 1 \right) \times \left(\sum_{(cd) \in \square} 4s_c^z s_d^z + (\bar{c}_{c\downarrow}^\dagger \bar{c}_{d\uparrow} + \bar{c}_{c\uparrow}^\dagger \bar{c}_{d\downarrow} + \bar{c}_{c\downarrow} \bar{c}_{d\uparrow} + \bar{c}_{c\uparrow} \bar{c}_{d\downarrow} + h.c.) + 3 \right) \times \left(\sum_{(fg) \in \square} 4s_f^z s_g^z + (\bar{c}_{f\downarrow}^\dagger \bar{c}_{g\uparrow} + \bar{c}_{f\uparrow}^\dagger \bar{c}_{g\downarrow} + \bar{c}_{f\downarrow} \bar{c}_{g\uparrow} + \bar{c}_{f\uparrow} \bar{c}_{g\downarrow} + h.c.) + 4 \right) \right]. \quad (24)$$

The states given by Eq.(22) are exact ground states of this short-range Hamiltonian. The frustrated kinetic hopping parameters and competing nearest neighbor and next nearest neighbor antiferromagnetic interactions physically adhere to the resulting ground state. The precise form of the diagonal formation of singlet dimers and high degeneracy of these states are non-trivial. Similar degeneracies appear in many other systems [18] where effective dimensional reduction occurs. By extending the correlator of Eq.(14) to the $S = 1/2$ arena by means of the transformations of [15] and noting that $\exp[2\pi i s_i^z] = \exp[i\pi \bar{n}_i]$ at all lattice sites i , we find a counterpart to the Ogata-Shiba correlators [17] for two leg ladders

$$\langle \tilde{\chi}_P | s_{1,i}^z s_{2,i}^z \rangle \exp[i\pi \sum_{k=i+1}^{j-1} (\bar{n}_{1,k} + \bar{n}_{2,k})] s_{1,j}^z s_{2,j}^z |\tilde{\chi}_P\rangle = \frac{1}{36}. \quad (25)$$

Here, $|\tilde{\chi}_P\rangle = \prod_{ij \in P} |\chi_{ij}\rangle$.

Other ground states can easily be proven along the same lines. As the fully ferromagnetic states $S_i^z = +1$ for all i or $S_i^z = -1$ are ground states of the spin $S = 1$ Hamiltonian

$$H = - \sum_{\square} \vec{S}_\square^2, \quad (26)$$

the maximally occupied (by hard core bosons) fully polarized ferromagnetic states $s_i^z = 1/2$ ($n_i = 1$) for all i or $s_i^z = -1/2$ ($n_i = 1$) at all sites i are ground states of the $s = 1/2$ Bose extended $t - J_z$ model

$$H = - \sum_{\square} \left[\sum_{(ab) \in \square} 4s_a^z s_b^z + (\bar{c}_{a\downarrow}^\dagger \bar{c}_{b\uparrow} + \bar{c}_{a\uparrow}^\dagger \bar{c}_{b\downarrow} + \bar{c}_{a\downarrow} \bar{c}_{b\uparrow} + \bar{c}_{a\uparrow} \bar{c}_{b\downarrow} + h.c.) \right]. \quad (27)$$




FIG. 2: (Color online.) From [10]. Highly regular ground states on the checkerboard and pyrochlore lattices. The ovals denote singlet dimer states. The arrows denote the representations of these dimer states within the six-vertex model (see text). On each plaquette (tetrahedron) the dimer connects the bases of the two incoming arrows.

III. OTHER LATTICES

We may repeat our derivations above to derive the ground states on any other lattice in which four fermions in any given basic unit interact with each other according to Eq.(20). One such example is afforded by the three dimensional pyrochlore lattice and its dimensional checkerboard rendition. The pyrochlore lattice is composed of corner sharing tetrahedra such that each vertex is common to two 4-site units (see the right hand side of Fig.(2)). The geometry is rather common in transition metal and rare earth oxide systems, specifically in the pyrochlore (the origin of the name) and spinel structures. The checkerboard lattice (see the left hand side of Fig.(2)) is a 2d projection of the pyrochlore structure: the crossed plaquettes in the left hand side of Fig.(2) represent the basic tetrahedral units.

On the pyrochlore/checkerboard lattices the open structure of tetrahedra/crossed plaquettes creates, by comparison to the square lattice, a less constrained system, and the number of dimer configurations in the ground-state manifold is strongly

enhanced, the still more massive degeneracy implying similarly exotic physics in this case.

In addition to the simple coverings shown in Fig. 2, a far richer variety of states exists. A detailed study of dimer states on the pyrochlore and checkerboard lattices is provided in [10]. A mapping onto the spin-ice problem provides a useful classification of these ground states. Here, the states with one dimer per tetrahedral unit defined by Eq. (8) map exactly to the spin-ice problem [19, 20] where, motivated by the structure of H atoms in solid water [21, 22], two of the sites of any elementary unit (a tetrahedron) are associated with an ingoing arrow pointing towards the center of the tetrahedron and two sites lie on arrows pointing outwards. We label the two sites belonging to a singlet dimer $|S_{ij}\rangle$ in a given tetrahedron $\square$ by two incoming arrows from sites i and j to the center of the unit $\square \equiv ijkl$. In this fashion it is clear that the system is mapped to a set of continuous directed lines such that each tetrahedral unit has exactly two incoming and two outgoing arrows relative to its center. An example of this mapping is illustrated in Fig. 2. The mapping is one-to-one: any spin-ice configuration determines a unique singlet-covering state with one dimer per tetrahedral unit and vice versa. The algebraic correlations in the spin-ice model dictate an algebraic decay of correlations in the $d = 2, 3$ dimensional $S = 1/2$ Fermi problems of Eq.(20) on the checkerboard and pyrochlore lattices: $\langle T_{ab} T_{mn} \rangle \sim |r|^{-d}$ for large separations $|r|$. Here, $|r|$ denotes the distance between the two site pairs $((ab)$ and $(mn))$ [10], [23], [24].

IV. CONCLUSION

In conclusion, we constructed spin $S = 1/2$ extended $t - J_z$ models (both fermionic and hard core bosonic) which harbor easily provable exact ground states. We illustrated how this method can be used to engineer non-uniform diagonal dimer state and other configurations. Applying similar ideas, many other solvable systems can be constructed on other lattices.

This work was partially sponsored by the Swedish Research Council.

-
- [1] J. G. Bednorz and K. A. Müller, Z. Phys. B – Condens. Matter **64**, 189 (1986)
 - [2] S. R. White and D. J. Scalapino, cond-mat/9907375, Phys. Rev. Lett. **80**, 1272 (1998);
 - [3] J. Zaanen and O. Gunnarsson, Phys. Rev. B **40**, 7391 (1989); K. Machida, Physica C **158**, 192 (1989)
 - [4] V. J. Emery and S. A. Kivelson, Physica C **209**, 597 (1993)
 - [5] Assa Auerbach, *Interacting Electrons and Quantum Magnetism* (Graduate Texts in Contemporary Physics), Springer-Verlag (1994)
 - [6] I. Affleck, T. Kennedy, E.H. Lieb and H. Tasaki, Commun. Math. Phys. **115**, 477 (1988); Phys. Rev. Lett. **59**, 799 (1987)
 - [7] D. J. Klein, J. Phys. A Math. Gen. **15**, 661 (1982).
 - [8] J. T. Chayes, L. Chayes, and S. A. Kivelson, Commun. Math. Phys. **123**, 53 (1989).
 - [9] C. D. Batista and S. A. Trugman, Phys. Rev. Lett. **93**, 217202 (2004).
 - [10] Z. Nussinov, C. D. Batista, B. Normand, and S. A. Trugman, cond-mat/0602528
 - [11] Z. Nussinov, cond-mat/0606075
 - [12] "Quasi-exactly solvable" here means that we can exactly determine all ground states yet do not solve the finite temperature problem.
 - [13] M. E. Fisher, Phys. Rev. **124**, 1664 (1961); P. W. Kasteleyn, Physica **27**, 1209 (1961)
 - [14] S. Todo, M. Matsumoto, C. Yasuda, H. Takayama, Phys. Rev. B **64**, 224412 (2001)
 - [15] C.D. Batista and G. Ortiz, Condensed Matter Theories Vol. **16**,

- ed. S. Hernandez and J. Clark (2001), 1
- [16] E. Fradkin, Phys. Rev. Lett., **63**, 322 (1989); D. Eliezer and G. W. Semenoff, Phys. Lett. B **286**, 118 (1992)
 - [17] M. Ogata and H. Shiba, Phys. Rev. B **41**, 2326 (1990); H. V. Kruis, I. McCulloch, Z. Nussinov, J. Zaanen, Phys. Rev. B **70**, 075109 (2004)
 - [18] C. D. Batista and Z. Nussinov, cond-mat/0410599, Phys. Rev. B, To appear
 - [19] S. T. Bramwell and M. J. P. Gingras, Science **294**, 1495 (2001), and references therein.
 - [20] P. W. Anderson, Phys. Rev. **102**, 1008 (1956).
 - [21] D. Bernal and R. H. Fowler, J. Chem. Phys. **1**, 515 (1933).
 - [22] L. Pauling, *The Nature of the Chemical Bond* (Cornell University Press, Ithaca, 1939).
 - [23] R. J. Baxter, *Exactly solved Models in Statistical Mechanics* (Academic Press, London, 1982).
 - [24] E. H. Lieb, Phys. Rev. **162**, 162 (1967).